

Interplay between heartbeat oscillations and wind outflow in microquasar IGR J17091-3624

Agnieszka Janiuk¹, Mikolaj Grzedzielski¹, Fiamma Capitanio², and Stefano Bianchi³

¹ Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotnikow 32/46, 02-668 Warsaw, Poland
e-mail: agnes@cft.edu.pl

² INAF – Istituto di Astrofisica e Planetologia Spaziali, via del Fosso del Cavaliere 100, 00133 Rome, Italy

³ Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, via della Vasca Navale 84, 00146 Roma, Italy

Received 16 September 2014 / Accepted 17 November 2014

ABSTRACT

Aims. During the bright outburst in 2011, the black hole candidate IGR J17091-3624 exhibited strong quasi-periodic flare-like events (on timescales of tens of seconds) in some characteristic states, the so-called heartbeat state. From the theoretical point of view, these oscillations may be modeled by the process of accretion disk instability, driven by the dominant radiation pressure and enhanced heating of the plasma. Although the mean accretion rate in this source is probably below the Eddington limit, the oscillations will still have large amplitudes. As the observations show, the source can exhibit strong wind outflow during the soft state. This wind may help to partially or even completely stabilize the heartbeat.

Methods. Using our hydrodynamical code GLADIS, we modeled the evolution of an accretion disk responsible for X-ray emission of the source. We accounted for a variable wind outflow from the disk surface. We examined the data archive from the *Chandra* and *XMM-Newton* satellites to find the observed limitations on the wind physical properties, such as its velocity and ionization state. We also investigated the long-term evolution of this source, which lasted over about 600 days of observations, using the data collected by the *Swift* and RXTE satellites. During this long period, the oscillations pattern and the observable wind properties changed systematically.

Results. We found that this source probably exhibits observable outbursts of appropriate timescales and amplitudes as a result of the disk instability. Our model requires a substantial wind component to explain the proper variability pattern, and even complete suppression of flares in some states. The wind mass-loss rate extracted from the data agrees quantitatively well with our scenario.

Key words. accretion, accretion disks – black hole physics – instabilities – X-rays: binaries

1. Introduction

The microquasar IGR J17091-3624 (hereafter IGR J17091) is a moderately bright transient X-ray binary (peak flux level at ~ 20 mCrab in the range 20–100 keV) discovered by INTEGRAL/IBIS in 2003 (Kuulkers et al. 2003) and classified as a black hole candidate (BHC) for its spectral and timing properties (Lutovinov & Revnitsev 2003; Capitanio et al. 2005; Lutovinov et al. 2005). After the INTEGRAL discovery, IGR J17091 was searched in the archival data of previous X-ray missions. The data of both TTM-KVANT (Revnitsev et al. 2003) and the *BeppoSAX* Wide Field Camera (WFC; in't Zand et al. (2003)) archives show that there were previous outbursts detected in 1994, 1996, and 2001.

The refined position of IGR J17091 provided by Kennea & Capitanio (2007) ruled out the tentative radio counterpart previously proposed for the source (Rupen et al. 2003; Pandey et al. 2006). The hypothesis was supported by reanalyzing the archival radio observations that identified a faint transient radio source (sub-mJy level at 5 GHz) that showed a flux increase immediately after the 2003 outburst. The inverted spectrum provides a signature of a compact radio jet (Capitanio et al. 2009) that is consistent with the low/hard spectral state (LHS) observed by INTEGRAL in the same period (Capitanio et al. 2005).

At the end of January 2011, the *Swift*/BAT hard X-ray transient monitor reported a renewed activity from IGR J17091. A

long monitoring campaign was then carried out with *Swift*/XRT, starting on February 28. Moreover, the source was extensively observed by INTEGRAL and RXTE.

The 2011 outburst was by far the brightest outburst ever observed from IGR J17091, and the source flux increased up to 120 mCrab in the range 2–10 keV (Capitanio et al. 2012). Follow-up radio observations carried out with the ATCA telescope measured a flat spectrum (Corbel et al. 2011; Rodriguez et al. 2011a; Torres et al. 2011) associated with self-absorbed compact jets. Later on, Rodriguez et al. (2011a) also reported on the detection of a discrete jet-ejection event that is typically observed when a BHC undergoes the transition from its hard-intermediate state to the soft-intermediate state.

The RXTE simultaneous observation campaign revealed several timing features, such as a 0.1 Hz quasi-periodic oscillation (QPO), increasing in frequency with the source flux and spectral softening (Rodriguez et al. 2011b; Shaposhnikov 2011), or a ~ 10 mHz QPO (Altamirano et al. 2011c). Moreover, RXTE/PCA data showed a continuous progression of quasi-periodic flare-like events occurring at a rate of between 25 and 30 mHz (Altamirano et al. 2011c). This type of variability resembles the heartbeat variation observed in the BH binary GRS 1915+105. Altamirano et al. (2011b) reported a detailed study of the behavior of the flare-like events of IGR J17091 during the first 180 days of the outburst. This study classified the different types of flares with the same scheme as used by

Table 1. *XMM-Newton* and *Chandra* observations log and spectral and timing properties.

Instrument	ID	Date	Exposure	Flux (2–10 keV)	Source state	Wind	Radio det.	T_{in}	Γ
–	–	–	ks	ergs s ⁻¹ cm ⁻²	–	–	–	keV	–
XMM	0677980201	2011-03-27	1	1.7×10^{-9}	heartbeat	no	yes ¹	1.24 ± 0.02	2.3 ± 0.1
<i>Chandra</i>	12405	2011-08-01	31	1.6×10^{-9}	heartbeat	no	no	1.26 ± 0.01	1.9 ± 0.1
<i>Chandra</i>	12406	2011-11-06	27	1.9×10^{-9}	soft-quiet	yes	no	1.7 ± 0.1	2.1 ± 0.3
XMM	070038130	2012-09-29	41	0.2×10^{-9}	hard	no	–	0.6 ± 0.1	1.32 ± 0.04^2

Notes. ⁽¹⁾ Seven days before. ⁽²⁾ Cutoff at 100_{-30}^{+50} keV.

1 Belloni et al. (2000) for GRS 1915+105, which have been clas-
2 sified into 14 variability classes.

3 In addition to all these similarities between GRS 1915+105
4 and IGR J17091 (see e.g. Capitanio et al. 2012; Pahari et al.
5 2014, for details), a particularly striking difference is the X-ray
6 flux intensity during the flare-like events. This fact cannot
7 be easily explained because unlike for GRS 1915+105, for
8 IGR J17091 we do not know the properties of the binary sys-
9 tem such as the distance, the inclination angle, BH mass, or
10 the spin. The GRS 1915+105 flux emission during the heart-
11 beat states reaches the Eddington luminosity (Done et al. 2004)
12 and is believed to be related to disk oscillations. As Altamirano
13 et al. (2011a) assumed, if the heartbeat oscillations seen from
14 IGRJ17091 are interpreted as being due to the same mechanism
15 as in GRS 1915+105, then the apparent faintness of IGR J17091
16 remains unexplained unless a huge distance or an extremely
17 low BH mass is considered (lower than $3 M_{\odot}$). However, the
18 subsequent results that estimate the distance and the mass of
19 IGR J17091 did not confirm this hypothesis.

20 Rebusco et al. (2012) estimated by studying the high-
21 frequency QPOs of the source (see also Altamirano & Belloni
22 2012) a BH mass of $6 M_{\odot}$ for which the distance extrapolated
23 from Altamirano et al. (2011a) should be >22 kpc, which would
24 mean that it lies approximately just outside the Galaxy. On the
25 other hand, Rodriguez et al. (2011a) calculated that the distance
26 range is 11–17 kpc for a BH mass of $10 M_{\odot}$, considering the
27 transition luminosity between the hard state and the soft state
28 (Yu & Yan 2009). Finally, the optical and near-infrared coun-
29 terparts have been identified by Torres et al. (2011) during the
30 2011 outburst, while the results from spectral analysis indicate
31 that the source is viewed at high inclination angle (King et al.
32 2012; Capitanio et al. 2012; Rao & Vadawale 2012). Several
33 hypotheses have been proposed to explain the faintness of the
34 IGR J17091 heartbeats, such as the spectral deformation ef-
35 fects due to a high inclination angle that is favored by the data
36 (Capitanio et al. 2012), or a low or even retrograde spin (Rao &
37 Vadawale 2012).

38 Another peculiarity of this source is the particularly fast and
39 ionized wind observed during the soft spectral state (King et al.
40 2012). King et al. (2012) reported that *Chandra* observations re-
41 veal two absorption lines at about 6.91 and 7.32 keV. These two
42 lines can be associated with blueshifted He-like Fe XXV and
43 Fe XXVI, with velocities of 9000 and 15000 km/s. This pro-
44 jected outflow velocity is an order of magnitude higher than pre-
45 viously observed in stellar-mass black holes.

46 In this paper, we explain the heartbeat oscillations in
47 IGR J17091 as due to the intrinsic oscillations of an accretion
48 disk that is subject to a thermal-viscous instability. As recently
49 shown in Janiuk & Czerny (2011), such oscillations do not re-
50 quire a disk luminosity equal to the Eddington rate, and the disk
51 instability may also occur for lower luminosities, on the order
52 of 0.1 Eddington (the extension of the disk part that is unsta-
53 ble as a result of the dominant radiation pressure decreases for

low accretion rates). As we demonstrated before (Janiuk et al.
2002), the wind ejection from the inner parts of an accretion disk
may provide one possible mechanisms to partially stabilize these
oscillations and affect the variability amplitudes. Therefore, we
also investigated the relationship between the wind and the evo-
lution pattern of the heartbeat oscillations.

Using our hydrodynamical code, we model the evolution
of an accretion disk responsible for the X-ray emission of
the source. We account for a variable wind outflow from the
disk surface and follow the evolution of the oscillation phase
thanks to the extensive observations campaigns performed by
Swift and RXTE during the 2011 outburst of the source. We
finally compare the theoretical predictions with the constraints
found with the data analysis of four observations performed by
XMM-Newton and *Chandra* satellites.

2. Observations

2.1. Data analysis

IGRJ17091 was observed by *Chandra* and *XMM-Newton* during
the 2011 outburst for a total of four pointings (two pointings for
each telescope), as reported in Table 1.

The two *Chandra* observations were carried out with the
Advanced CCD Imaging Spectrometer (ACIS: Garmire et al.
2003) with the High-Energy Transmission Grating Spectrometer
(HETGS: Canizares et al. 2005) in the focal plane, and are pre-
sented in King et al. (2012). Data were reduced with the *Chandra*
Interactive Analysis of Observations (CIAO: Fruscione et al.
2006) 4.6 and the *Chandra* Calibration Data Base (CALDB)
4.6.1.1 software, adopting standard procedures. First-order HEG
spectra were extracted for the source and the background for a
total exposure time of 31 and 27 ks for the two observations. Fits
were performed on the unbinned spectra, using the Cash (1976)
statistics, to take advantage of the high spectral resolution of the
gratings.

The *XMM-Newton* observations were operated in burst and
timing mode with the European Photon Imaging Camera (EPIC)
CCD pn camera (Strüder et al. 2001). Data were reduced with
SAS 13.5.0, and the screening for intervals of flaring particle
background was made consistently with the choice of extraction
radii, in an iterative process based on the procedure to maximize
the signal-to-noise ratio described in detail by Piconcelli et al.
(2004) in their Appendix A. As a result, we extracted the source
spectrum from a box region 15 (17) RAWX pixels wide, cen-
tered on the source for observation 0677980201 (070038130),
while the background spectrum was extracted from RAWX
columns 2–9 and 57–63. The resulting good exposure times are 1
and 46 ks. Finally, spectra were binned to oversample the instru-
mental resolution by at least a factor of 3 and to have no fewer
than 30 counts in each background-subtracted spectral channel.
The latter requirement allowed us to use χ^2 statistics.




Fig. 1. X-ray light curve of the source IGR J17091 as observed by *Swift*/XRT. The observation ID is 31921041.

The source was also observed up to 200 keV with the γ -ray telescopes, IBIS (Ubertini et al. 2003), onboard the INTEGRAL satellite, simultaneously with *Chandra* and *XMM-Newton*. The INTEGRAL/IBIS data were analyzed using the latest release of the standard Offline Scientific Analysis, OSA, version 10.0, and the latest response matrices available. The spectral analysis was focused on ISGRI, the IBIS low-energy detector (Lebrun et al. 2003). The ISGRI spectra were extracted in the 20–200 keV energy range. A systematic error of 2% was taken into account, as reported, for example, by Jourdain et al. (2008). To collect enough counts in a single spectrum, we summed contiguous IBIS spectra that showed the same shape.

In the following, errors and upper limits correspond to the 90 per cent confidence level for one interesting parameter ($\Delta\chi^2 = 2.71$), unless otherwise stated. The data were analyzed with the HEASOFT package version v.6.15.1.

2.2. Time variability

To determine the timing behavior of the source and the presence or absence of periodical flare-like events associated with the heartbeat state during a period that includes the four *XMM-Newton* and *Chandra* observations (April 2011–October 2012), we analyzed the data collected by *Swift*/XRT and RXTE/PCA. We extracted the *Swift*/XRT 0.2–10 keV light curve using the standard XRT pipeline, while for RXTE/PCA data we used the sdt1 mode light curves.

Figure 1 shows an example of the time-variability pattern characteristic for the heartbeat state of the source. The light curve was extracted from *Swift*/XRT data observed on 17 April 2011 (ID: 00031921041, exposure: 1 ks). The observation was taken in window-timing mode to avoid pile-up effects. Typically, IGR J17091 shows a heartbeat variability with a period of a few tens of seconds, as previously reported by Altamirano et al. (2011b) and Capitanio et al. (2012).

2.3. Wind and heartbeat detection

Based on extensive *Swift* and RXTE observations, we report in Fig. 2 the periods in which the periodical flare-like events have been detected, superimposed on the 15–50 keV *Swift*/BAT light curve of IGR J17091.




Fig. 2. *Swift*/BAT 15–50 keV light curve superimposed on a time-line sketch created from the *Swift*/XRT, RXTE/PCA, *Chandra*, and XMM data analysis, showing the anticorrelation between the wind and heartbeat in IGR J17091. The second *Chandra* observation (second green arrow) shows a fast, ionized wind, while the first *XMM-Newton* observation (first red arrows) and the first *Chandra* observation (first green arrow) lie in the heartbeat zone and do not show any detectable wind outflow. During the last *XMM-Newton* observation (second red arrow), the source is in a hard state. The parts with no flare are marked with red-oblique-dashed lines. Finally the green-oblique-dashed lines individuate periods without observations.

Two simultaneous observations made by *EVLA* at the time of each *Chandra* pointing did not detect any radio emission (King et al. 2012), confirming that if there is an outflow, it is probably only due to the wind ejection. However, an *ATCA* observation detected a radio counterpart of IGR J17091 seven days before the first *XMM-Newton* observation (Corbel & Tzioumis 2011). This means that a contribution of the jet emission to the outflow cannot be excluded a priori in this case.

XMM-IBIS and *Chandra-IBIS* joined spectra were modeled with the simplest possible model: an absorbed multicolor disk black body (Shakura & Sunyaev 1973) plus a power-law. Table 1 summarizes the results of the spectral fitting.

The first *Chandra* observation does not show any significant absorption line in the spectrum that could be associated with a wind outflow (King et al. 2012). The same is true for the first *XMM-Newton* observation, with the exception of a tentative detection of a feature at ≈ 7.1 keV (see Reis et al. 2012)¹. A prominent heartbeat is present in the light curves of both observations. On the other hand, the second *Chandra* observation shows a strong and fast wind, as reported by King et al. (2012), but no detection of any strong periodical flare-like events. Finally, the spectral analysis of the second *XMM-Newton* observation indicates that the source is in a hard state (see Table 1): our analysis does not show any absorption line, in agreement with the general behavior of GBHs in this state (Ponti et al. 2012).

In the next section, we study in detail the two *Chandra* observations for which it is indicated that the outflow can only be ascribed to the wind ejection; the high-resolution grating spectra allow us to characterize or exclude these winds.

2.4. Photoionization modeling of the data

To characterize the wind in the *Chandra* observations, we prepared two tables with the photoionization code Cloudy C13.02 (last described in Ferland et al. 2013). We adopted the two spectral energy distributions (SEDs) described in the previous

¹ Our analysis confirms this isolated absorption feature, detected with a nominal confidence level of $\approx 3\sigma$.




Fig. 3. *Chandra* ACIS-S HETG spectra for observations 12406 (upper panel) and 12405 (lower panel) in the 6–7.5 keV energy range. The wind components observed in observation 12406 are completely undetectable in observation 12405 if we assume that the wind density is a factor of 10 lower. See text for details.

section, and, following King et al. (2012), a turbulence velocity $v_{\text{turb}} = 1000 \text{ km s}^{-1}$ and an overabundance of iron by a factor of two (the other chemical abundances are as in Table 7.1 of the Cloudy documentation).

We first fitted the second *Chandra* observation with a continuum model resembling that presented in King et al. (2012), that is, a disk black-body and a power law. Two photoionized components are required to fit the absorption residuals around 6–7 keV: one with an outflow velocity $v_1 = 9700^{+800}_{-700} \text{ km s}^{-1}$, $\log \xi_1 = 3.4^{+0.2}_{-0.3}$, and $\log N_{\text{H}1} = 22.1^{+0.2}_{-0.4}$; the other with an outflow velocity $v_2 = 15700 \pm 600 \text{ km s}^{-1}$, $\log \xi_2 = 3.8^{+0.2}_{-0.1}$, and $\log N_{\text{H}2} = 22.5 \pm 0.3$. We note here that these values perfectly agree with those reported by King et al. (2012). The spectrum and best-fit model are shown in Fig. 3.

From the best-fit values of the ionization parameter $\xi = L_{\text{ion}}/nr^2$, assuming the ionizing (1–1000 Ryd) luminosity as in the input SED of $L_{\text{ion}} = 3.7 \times 10^{37} \text{ erg s}^{-1}$, we derive $n_1 r_1^2 \approx 1.5 \times 10^{34}$ and $n_2 r_2^2 \approx 5.8 \times 10^{33} \text{ cm}^{-1}$ for the two wind components. The corresponding (spherical) mass outflow rates would be

$$\dot{M}_{\text{wind}} \approx 1.23 m_p f v \Omega n r^2, \quad (1)$$

where m_p is the proton mass (the correction factor of 1.23 comes from assuming cosmic elemental abundances), Ω is the covering factor, and f the filling factor. The absence of emission lines suggests $\Omega/4\pi \approx 0.5$ (King et al. 2012), so our estimated mass outflow rates are

$$\dot{M}_{\text{wind}1} \approx 1.8 \times 10^{20} f_1 \text{ g s}^{-1} \quad (2)$$

$$\dot{M}_{\text{wind}2} \approx 1.2 \times 10^{20} f_2 \text{ g s}^{-1}. \quad (3)$$

The filling factors f_1 and f_2 of the wind components are more difficult to estimate. It is reasonable to assume that the extension of the wind is similar to its launching radius, $r \approx \Delta r$, so that the observed absorption column densities are related to the filling factors, which means that $N_{\text{H}} = fnr$. To break the degeneracy between n and r , we assumed that the outflow velocity of the wind is on the order of the escape velocity at its launching radius. This leads to $r_1 \approx 1900 r_g$ and $r_2 \approx 760 r_g$, corresponding to $n_1 \approx 5.1 \times 10^{15} \text{ cm}^{-3}$ and $n_2 \approx 1.3 \times 10^{16} \text{ cm}^{-3}$. The resulting

filling factors, $f_1 \approx 0.0015$ and $f_2 \approx 0.0037$, give $\dot{M}_{\text{wind}1} \approx 2.7 \times 10^{17} \text{ g s}^{-1}$ and $\dot{M}_{\text{wind}2} \approx 4.2 \times 10^{17} \text{ g s}^{-1}$.

The first *Chandra* observation does not show any absorption feature. If we assume that the density of the wind is lower by a factor of 10 with respect to the density characterizing the observed wind in the second observation, we would expect ξ and N_{H} to re-scale linearly (but we also take into account the small difference in the SED, and, therefore, in L_{ion}). As shown in Fig. 3, the two wind components would be undetectable in our data if they had these parameters. Assuming that all the other properties of the wind remain the same, we obtain lower mass outflow rates of $\dot{M}_{\text{wind}1} \approx 2.5 \times 10^{16} \text{ g s}^{-1}$ and $\dot{M}_{\text{wind}2} \approx 3.9 \times 10^{16} \text{ g s}^{-1}$.

3. Model of the radiation pressure instability

The outbursts of accretion disks may be induced by the two main types of instabilities that lead to the thermal-viscous oscillations. These are the radiation pressure instability and the partial hydrogen ionization instability, both known for over 40 years in theoretical astrophysics. For a detailed discussion of these instabilities we refer to Janiuk & Czerny (2011) and to the literature quoted therein. We here studied the accretion disk instability induced by the dominant radiation pressure and modeled it to explain the behavior of the microquasar IGR J17091.

The stationary, thin accretion disk model in classical theory is based on an α prescription for the viscous energy dissipation. In the α -model we assume the nonzero component $T_{r\phi}$ of the stress tensor to be proportional to the total pressure. The latter includes the radiation pressure component, which scales with temperature as T^4 and blows up in hot disks for high accretion rates. This in turn affects the heating and cooling balance between the energy dissipation and radiative losses.

This balance, assuming hydrostatic equilibrium, is calculated numerically with a closing equation for the locally dissipated flux of energy given the black hole mass and global accretion rate. The local solutions may be conveniently plotted on the so-called stability curve. This S-shaped scheme represents an annulus in the accretion disk, with temperature and surface density determined by the accretion rate. If the accretion rate is low, the annulus of the disk is dominated by the gas pressure and stable. For higher accretion rates, the annulus is dominated by radiation pressure and unstable. An upper stable branch of this curve appears for even higher accretion rates, close to the Eddington limit, and is due to efficient advective cooling. The higher the global accretion rate, the more annuli of the disk will be affected by this instability. In consequence, for a larger size of the instability zone, the disk will undergo stronger amplitude oscillations. Below a critical accretion rate, the instability zone completely vanishes and the whole disk is dominated by gas pressure and hence stable. We determined this rate to be about 3.6 per cent Eddington.

3.1. Physics of the model

3.1.1. Equations

We assumed nondiagonal terms in stress tensor in cylindrical coordinates proportional to the total pressure with a constant viscosity α , as introduced by Shakura & Sunyaev (1973),

$$T_{r\phi} = -\alpha P, \quad (4)$$

where P is the pressure.

1 If we consider a thin disk whose energy loss is only con-
2 nected with horizontal flows, local energy loss in the disk per
3 unit volume per unit time is given by

$$\frac{\partial F}{\partial z} = T_{r\phi} \frac{d\Omega}{dr}, \quad (5)$$

4 where v_ϕ is the angular (Keplerian) velocity in the disk

$$\Omega_K = \sqrt{\frac{GM}{r^3}}. \quad (6)$$

5 It is convenient to express the quantities in terms of Eddington
6 units, and the dimensionless accretion rate is

$$\dot{m} = \frac{\eta \dot{M} c^2}{L_{\text{Edd}}} = \frac{\dot{M}}{\dot{M}_{\text{Edd}}}. \quad (7)$$

7 Here η is the efficiency of accretion, equal to 1/16 in the pseudo-
8 Newtonian approximation. According to Eqs. (4) and (5), the
9 final formula for the viscous energy dissipation is

$$\frac{\partial F_{\text{tot}}}{\partial z} = \alpha P \frac{d\Omega}{dr}. \quad (8)$$

10 The equilibrium pressure consists of gas and radiation pressure,

$$P = P_{\text{gas}} + P_{\text{rad}}, \quad (9)$$

11 where we assume that gas is an ideal gas of protons of mass m_p ,

$$P_{\text{gas}} = \frac{\rho k_B T}{m_p}, \quad (10)$$

12 and the radiation pressure is given by

$$P_{\text{rad}} = \frac{4\sigma_B T^4}{3c}. \quad (11)$$

13 We consider here the model of a vertically averaged disk (effec-
14 tively, a 1.5-dimensional model, as the motion in the angular di-
15 rection is also accounted for). In the simplest possible stationary
16 case, we close the equations by an energy conservation equation
17

$$F_{\text{tot}} = Q_+ = Q_-, \quad (12)$$

18 where the total flux locally emitted from the surface of the ac-
19 cretion disk is given by the global parameters

$$F_{\text{tot}} = \frac{3GM\dot{M}}{8\pi r^3} f(r), \quad (13)$$

20 where $f(r)$ is given by the boundary condition on the inner edge
21 of accretion disk. The vertically averaged viscous heating rate in
22 the disk is

$$Q_+ = \frac{3}{2} \alpha PH \Omega_K, \quad (14)$$

23 and the radiative cooling rate per unit time per surface unit is

$$Q_- = \frac{4\sigma_B T^4}{3\kappa \Sigma}, \quad (15)$$

24 where κ is the electron scattering opacity, which is equal
25 to $0.34 \text{ cm}^{-2} \text{ g}^{-1}$.

26 The advective cooling is included in the stationary model as

$$Q_- = F_{\text{tot}}(1 - f_{\text{adv}}), \quad (16)$$

where f_{adv} is defined as follows (Muchotrzeb & Paczynski 1982;
Abramowicz et al. 1988; Janiuk et al. 2002):

$$f_{\text{adv}} = \frac{-2rP}{3\rho GM f(r)} q_{\text{adv}}, \quad (17)$$

and q_{adv} is given by

$$q_{\text{adv}} = (12 - 10.5\beta) \frac{d \ln T}{d \ln r} - (4 - 3\beta) \frac{d \ln \rho}{d \ln r}, \quad (18)$$

where $\beta = P_{\text{gas}}/P$. The fraction q_{adv} in stationary model is as-
sumed to be a constant on the order of unity.

3.1.2. Time-dependent 1.5D hydrodynamics

In the time-dependent model we solve the full set of equations of
hydrodynamics. We consider a thin-disk model, so in the conser-
vation of mass (continuity) equation we neglect the z -behavior
of density and velocity fields:

$$\frac{\partial(\Sigma v_r)}{\partial r} + r \frac{\partial \Sigma}{\partial t} = 0. \quad (19)$$

The mass conservation equation is therefore

$$\frac{\partial \Sigma}{\partial t} = \frac{1}{2\pi r} \frac{\partial \dot{M}}{\partial r}, \quad (20)$$

with

$$\dot{M} = -2\pi r \Sigma v_r. \quad (21)$$

The angular momentum conservation is included in the equation

$$\dot{M} \frac{d}{dr} (r^2 \Omega) = -\frac{\partial}{\partial r} (2\pi r^2 T_{r\phi}). \quad (22)$$

We define ν as the kinematic viscosity

$$T_{r\phi} = \alpha PH = \frac{3}{2} \Omega \nu \Sigma. \quad (23)$$

From the mass and angular momentum conservation Eq. (22) we
obtain the final formula on the evolution of the surface density
of disk:

$$\frac{\partial \Sigma}{\partial t} = \frac{1}{r} \frac{\partial}{\partial r} (3r^{1/2} \frac{\partial}{\partial r} (r^{1/2} \nu \Sigma)). \quad (24)$$

It has the form of a simple diffusion equation when we transform
the variables

$$\frac{\partial \Xi}{\partial t} = \frac{12}{y^2} \frac{\partial^2 \Xi}{\partial y^2} (\Xi \nu), \quad (25)$$

where $r = 2y^{1/2}$ and $\Xi = 2r^{1/2} \Sigma$. The radial velocity is given by

$$v_r = -\frac{3}{\Sigma} r^{-1/2} \frac{\partial}{\partial r} (\nu \Sigma r^{1/2}). \quad (26)$$

To close the set of the equations, we need the energy conserva-
tion equation. In the disk we may have radiative cooling as well
as convection, advection, and diffusion. In our model we neglect
the effects of convection and heat diffusion and assume the same
equation as in Janiuk et al. (2002),

$$\frac{\partial \ln T}{\partial t} + v_r \frac{\partial \ln T}{\partial r} = \frac{4 - 3\beta}{12 - 10.5\beta} \left(\frac{\partial \ln \Sigma}{\partial t} - \frac{\partial \ln H}{\partial t} + v_r \frac{\partial \ln \Sigma}{\partial r} \right) + \frac{Q_+ - Q_-}{(12 - 10.5\beta)PH}. \quad (27)$$

Table 2. Resulting heartbeat outbursts modeled with different grids.

N_p	Period	Error	Variance	Duration of outburst	Error	Variance	N_o
25	441.16	52.7	341.63	10.68	0.38	2.52	44
30	684	77.9	397.3	29.28	3.81	19.81	27
40	1199	47.6	184.4	57.06	2.81	11.25	16
50	1342	43.8	158.0	70.66	3.45	12.89	15
75	1381	21.1	76.0	76.67	2.11	7.89	14
100	1402	18.5	64.2	78.57	2.06	7.42	13

Notes. The first column, N_p , gives the number of grid points. The last column, N_o , gives the number of outbursts in the time interval of 20 ks. The other quantities are given in seconds.

3.2. Numerics

We used the code global accretion disk instability (GLADIS), whose basic framework was initially described by Janiuk et al. (2002). The code was subsequently developed and applied in a number of works to model the evolution of accretion disks in Galactic X-ray binaries and AGN (Janiuk & Czerny 2005; Czerny et al. 2009). The current version includes the parallelization through the message-passing interface method (MPI; Janiuk & Misra 2012) and allows for computations with a variable time-step down to a thermal timescale, adjusting to the speed of local changes of the disk structure.

3.2.1. Grid and regularity

First, we tested the code parameterized with the accretion rate equal to the Eddington rate, that is, $\dot{m} = 1$, and a central object mass of $M = 10 M_\odot$. The grid covers the area from the inner radius R_{in} equal to 3 Schwarzschild radii to external radius equal to 100 Schwarzschild radii.

In Table 2 we summarize the tests of our numerical grid. We computed the average value \bar{T} in our modeled light curve as

$$\bar{T} = \frac{1}{N} \sum_{i=1}^N T_i, \quad (28)$$

where $T_i = t_{i+1} - t_i$ are distances between $i + 1$ th and i th outburst peaks. The variance V_T of periods for each grid is

$$V_T = \sqrt{\frac{1}{N} \sum_{i=1}^N (T_i - \bar{T})^2}, \quad (29)$$

and errors are given as

$$E_T = \sqrt{\frac{1}{N(N-1)} \sum_{i=1}^N (T_i - \bar{T})^2}. \quad (30)$$

We note that a denser grid leads to a better regularity of outbursts. For a number of grid points equal to 100 the average period is equal to 1401.92 s, with variance of periods between outbursts equal to 64.1 s, which is less than 5 per cent. We conclude that outbursts are regular in our model and accurately describe the heartbeat effect.

3.3. Parameters

We adopted the global parameters for the accretion disk model to be the black hole mass of IGR J17091 equal to $6 M_\odot$, as estimated by Rebusco et al. (2012) and (arbitrary) viscosity parameter $\alpha = 0.1$. To determine the accretion rate, we determined the distance of IGR17091 using the same method as reported by

Rodriguez et al. (2011a) rescaled for $6 M_\odot$, which gives a distance range of $d = 8-13$ kpc. The absolute luminosity range based on this distance range is $L = 3.1 \times 10^{37} - 8.2 \times 10^{37} \text{ erg s}^{-1}$. For the black hole mass adopted here this gives a dimensionless accretion rate range of $\dot{m} = 0.041-0.108$, which is equal to $\dot{M} = 0.86-2.12 \times 10^{-8} M_\odot \text{ yr}^{-1}$ for this source.

3.4. Results

The radiation pressure instability model with the parameters given above leads to the pronounced, regular luminosity outbursts. Obviously, the heating proportional to the gas pressure results in a completely stable accretion disk, with a constant luminosity, regardless of the accretion rate. Moreover, if we assume the viscous heating given by the geometrical mean of the gas and radiation pressures, no outbursts occur. Therefore, to model the heartbeat states of IGR J17091, we need the total pressure prescription for the viscous torque (Eq. (23)).

3.4.1. Behavior of the disk during an outburst cycle

If there were no stabilizing mechanism, the radiation-pressure dominated disk would not survive. This is because in these parts of the disk the decreasing density leads to the temperature growth. In consequence, the local accretion rate increases and more material is transported inward. The disk annulus empties because of both the increasing accretion rate and decreasing density, so there is no self-regulation of the disk structure. However, the so-called slim-disk solution, where advection of energy provides an additional source of cooling in the highest accretion rate regime (close to the Eddington limit), acts as a stabilizing branch. Therefore, even if a large part of the disk is dominated by radiation pressure, advection of some part of energy allows the disk to survive and oscillate between the hot and cold states. This oscillating behavior leads to periodic changes of the disk luminosity, as shown, for instance, in Janiuk et al. (2000) for the black hole X-ray binary disks.

In Fig. 4 we show the distributions of density and temperature in the accretion disk in the peak of an outburst. The quantities are vertically averaged, while we also computed the disk thickness, as indicated on the y -axis. The ratio of H/r is always lower than about 0.1, which justifies our approach. In the outburst peak, the inner disk becomes flaring and its temperature rises, while the density drops by an order of magnitude.

Figure 5 shows an exemplary light-curve, with oscillations on a timescale of a few hundreds of seconds, as calculated for the black hole mass and accretion rate given in Sect. 3.3. The amplitudes of these outbursts, however, are much larger than those observed in the source's X-ray light-curves, and by over an order of magnitude exceed the luminosity changes detected in IGR J17091.




Fig. 4. Profiles of the disk density and temperature during an outburst in cgs units. The distribution is vertically averaged (note that the approximation of a geometrically thin disk also holds during the flare). Parameters: viscosity $\alpha = 0.1$, mean accretion rate $\dot{m} = 0.1$, and black hole mass $M = 6 M_{\odot}$. Distance is expressed in Schwarzschild radius units.




Fig. 5. Example of light curve modeled with an accretion disk instability, for $A = 0$ (without wind), a mean accretion rate $\dot{m} = 0.1$, and a black hole mass $M = 6 M_{\odot}$.

3.4.2. Partial stabilization of the disk outbursts as a result of wind outflow

As known before (e.g., Janiuk & Czerny 2005; see also Neilsen et al. 2012), the disk oscillations may be partially stabilized by the wind outflow and/or corona. Here we suggest that part of




Fig. 6. Dependence of the outburst amplitudes and periods on the wind strength A . Shaded regions mark observed amplitudes and timescales of the microquasar heartbeats. Different colors mark different values of the parameter, as indicated in the plot. Models were ran for \dot{m} between 0.04 and 0.1. In general, the lower the accretion rates, the shorter the outbursts, as long as the instability occurs.

the energy generated in the accretion process is used to accelerate the particles to eject the outflow from the disk surface. The reduced heating will result in smaller amplitude outbursts produced through the instability.

Following Janiuk et al. (2002), we corrected for the fraction of energy that is transferred to the disk surface (so in principle, we modified the conditions of the vertical energy transport that are behind the vertically integrated model). We adopted a function that assumes wind-launching power scales quadratically with the local accretion rate:

$$f_{\text{out}} = 1 - \frac{1}{(1 + Am^2)}. \quad (31)$$

Various other mathematical prescriptions were studied, for example, in Nayakshin et al. (2000), who found no qualitative importance on the details of this function. This scaling represents the physical nature of the process and simply means that the wind particles are ejected when the local disk luminosity instantaneously approaches the Eddington limit: $\dot{m} = \dot{M}/\dot{M}_{\text{Edd}} = L/L_{\text{Edd}}$. Here f_{out} is a dimensionless number in the range (0,1), and in our computations it depended on both radius and time via $\dot{M} \propto r\Sigma v_r$. This fraction modifies the local heating-to-cooling balance, which means that effectively f_{out} enters into the denominator in Eq. (??). The free constant $A \geq 0$ is our model parameter; it describes the strength of the wind and its mass-loss rate. Note that in the evolutionary calculations, we did not update the local density of the disk according to this mass loss because the latter is lower by many orders of magnitude.

We first phenomenologically determined the value of the wind strength parameter A , so that the resulting reduced outburst amplitudes with nonzero wind outflow reproduce the observed variability scale of the microquasar IGR J17091, like those shown in Fig. 1.

Figure 6 shows the dependence of outburst amplitudes on the changing wind strength parameter. Different symbols in this figure mark different values of A , and we show several results for each A , depending on the adopted mean accretion rate, \dot{m} , as taken in the range of values determined for microquasar IGR J17091. Values lower than $A = 10$ were excluded because




Fig. 7. Model light curve of the accretion disk. The luminosity changes are due to the radiation pressure instability. The wind strength is $A = 15$. Other parameters: black hole mass $M = 6 M_{\odot}$, viscosity $\alpha = 0.1$, accretion rate $\dot{M} = 2.12 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$ ($\dot{m} = 0.1$).

1 they yielded too large amplitudes and long outburst timescales
2 for any plausible accretion rate. On the other hand, values of
3 $A > 25$ yield too small amplitudes, and outbursts cease com-
4 pletely for $A = 100$.

5 Figure 7 shows an exemplary light curve from the model
6 with wind, which best represents the behavior of our source in
7 its heartbeat state. Since the accretion disk emission flux peaks
8 at about 10 Schwarzschild radii, the light curve shown here cor-
9 responds to the soft X-ray band for a stellar-mass accreting black
10 hole.

11 3.4.3. Average mass loss, density, and column density 12 of the wind

13 The mass-loss rate in the vertical direction is equal to the ra-
14 tio of the locally generated flux in the accretion disk, with a
15 fraction determined by f_{out} , to the energy change per particle,
16 $\dot{m}_z = F_{\text{tot}} f_{\text{out}} / (\Delta E / m_p)$. The local flux is given by Eq. (13)
17 of the standard accretion disk theory, and we assumed that this re-
18 lation also holds in the hydrodynamical computations. The en-
19 ergy change per particle is on the order of the virial energy,
20 $\Delta E = B k T_{\text{vir}} = B G M m_p / r$, with $B \sim 1$, so that expressing
21 the mass-loss rate in terms of local variables, in the units of
22 $[\text{g s}^{-1} \text{ cm}^{-2}]$, we have the final formula

$$\dot{m}_z(r) = B^{-1} \frac{3}{4} \frac{1}{r} \Sigma v_r f(r) f_{\text{out}}, \quad (32)$$

23 with f_{out} given by Eq. (31).

24 To obtain the total mass loss in the wind, we integrated \dot{m}_z
25 over the disk:

$$\dot{M}_{\text{wind}} = \int_{R_{\text{min}}}^{R_{\text{max}}} \dot{m}_z 4\pi r dr. \quad (33)$$

26 We assumed the outer radius of the accretion disk to be equal
27 to $4 \times 10^4 R_g$, which is the largest possible size of the disk
28 in IGR J17091, as estimated from the supposed parameters of
29 the binary. The smallest radius, R_{min} , could be located at the
30 marginally stable orbit around the black hole, that is, at $3 R_g$ for
31 a nonrotating hole, or anywhere above it. We therefore list sev-
32 eral results here for the smallest radii from which the unbound
33 wind is launched.

Table 3. Exemplary results of the disk/wind model.

R_{min}	$\dot{M}_{\text{wind}} [\text{g s}^{-1}]$	$\rho_{\text{wind}} [\text{g cm}^{-3}]$	$N_{\text{H}} [\text{cm}^{-2}]$
3	2.18×10^{17}	2.0×10^{-16}	5.66×10^{21}
70	8.84×10^{16}	3.9×10^{-16}	1.10×10^{22}
150	7.68×10^{16}	5.0×10^{-16}	1.07×10^{22}
350	6.04×10^{16}	7.1×10^{-16}	2.00×10^{22}
2000	3.06×10^{16}	7.2×10^{-16}	2.04×10^{22}

Notes. The parameters are the accretion rate $\dot{m} = 0.1$, wind strength $A = 15$, black hole mass $M = 6 M_{\odot}$, and wind extension $R_{\text{max}} = 40\,000 R_g$. The column density is calculated assuming $f_1 = 0.0015$.

The actual mass loss will be on this order, or somewhat
smaller, as the wind particles may be accelerated to obtain ki-
netic energy not necessarily equal to the virial energy, but rather
to have a velocity a few (2–3) times their escape velocity at the
radius r . Here we therefore calculate the upper limit for the wind
column density, depending on our model parameter A .

The wind mass-loss changes in time as a result of the out-
bursts. Averaging the solutions, we obtain the total mass loss
caused by the wind at the level of $\dot{M}_{\text{wind}} = 3 \times 10^{16} - 2 \times$
 10^{17} g s^{-1} .

We calculate the wind density at R_{max}

$$\rho_{\text{wind}} = \frac{\dot{M}_{\text{wind}}}{4\pi R_{\text{max}}^2 v_{\text{esc}}}, \quad (34)$$

where v_{esc} is the escape velocity at R_{min} . In our best-fit heart-
beat model, the density of the wind is equal to $\rho_{\text{wind}} = 2.0 \times$
 $10^{-16} - 7.2 \times 10^{-16} \text{ g cm}^{-3}$, depending on the wind-launching
radius.

Assuming that wind consists of protons, we calculate the col-
umn density of the (spherically symmetric) wind:

$$\int \rho_{\text{wind}}(\mathbf{r}) d\mathbf{r} = \rho_{\text{wind}}(R_{\text{max}}) R_{\text{max}}. \quad (35)$$

The observable column number density of particles is then be
given by $N_{\text{H}} = \rho_{\text{wind}} R_{\text{max}} (f m_p)^{-1}$, where f is the filling factor, in
principle unknown and constrained from the wind observations.
The resulting wind mass-loss rate and density for an assumed
wind strength parameter $A = 15$ and several values of the inner
radius at which the wind is launched are summarized in Table 3.

3.5. Comparison between the wind model and observations of IGR J17091

To reconcile the model with observations of the wind in
IGR J17091, we need to constrain the wind launching zone in
its radial extension to specify R_{min} and R_{max} . We also need to
determine the wind strength parameter A to either reproduce the
amplitude of the observed heartbeat oscillations or suppress them
completely. For radii below ~ 70 Schwarzschild radii, the depen-
dence of $\dot{m}_z(r)$ derived from Eq. (32) is a complicated function,
while at larger radii it scales with radius as a simple power law
with index -1.8 . Therefore, we fitted this relation and its nor-
malization to our best-fit model for the heartbeat oscillating disk,
that is, with $A = 15$. After simple integration, we obtain that

$$\dot{M}_{\text{wind}}(r, R_{\text{max}}) = 1.635 \times 10^{16} (R_{\text{max}}^{0.2} - r^{0.2}) [\text{g s}^{-1}], \quad (36)$$

where in the above the argument $r = R_{\text{min}} \geq 70$ and is given
in the units of $2GM/c^2$, and we already assumed the covering
factor of $\Omega = 2\pi$.

1 From this relation, we can constrain the extension of the
2 wind-launching zone, taking into account the radius r deter-
3 mined by the velocity of the wind equal to the escape velocity
4 at that radius, which can be constrained by the spectral analy-
5 sis (provided the wind is detectable in the data). Therefore,
6 if we adopted the outflow velocities to be the same as in the
7 second *Chandra* observation, then we can tentatively estimate
8 that for the first wind component with a mass-loss rate of about
9 $\dot{M}_{\text{wind1}} = 2.5 \times 10^{16} \text{ g s}^{-1}$, the extension of the launching zone
10 should be up to $R_{\text{max}} \approx 4900$ Schwarzschild radii. The second
11 wind component, with $\dot{M}_{\text{wind2}} = 3.9 \times 10^{16} \text{ g s}^{-1}$, should be
12 launched in the zone up to $R_{\text{max}} \approx 5900$.

13 Furthermore, we note the strong dependence of the wind
14 mass-loss rate on the strength parameter A .

15 With $A = 300$, a stable disk solution is found in our simula-
16 tions, while the mass-loss rate \dot{M}_{wind} is about ~ 12 times higher
17 than for our heartbeat model with $A = 15$. Such a strong wind
18 definitely stabilizes the heartbeat oscillations, while it should
19 give clear observable signatures in the *Chandra* spectra, such
20 as those presented in Fig. 3. Taking the values of the wind
21 velocities and mass-loss rates to be the same as those in the
22 first *Chandra* observation, we obtain observable wind-launching
23 zones of 950–4200 and 380–4700 Schwarzschild radii for the
24 first and second wind components.

25 4. Discussion

26 4.1. General picture for IGR J17091

27 The photoionization modeling of winds detected during the two
28 *Chandra* observations revealed a strong wind in the state without
29 heartbeat oscillations. We found two components of the wind,
30 characterized by different velocities and hence different launch-
31 ing radii and mass outflow rates.

32 In the heartbeat state, on the other hand, we suspect that
33 wind is also ejected from this source, albeit with a density lower
34 by about ten times and therefore undetectable from the absorp-
35 tion lines. The mass outflow rates that we determined from the
36 observation analysis is 2.7 and $4.2 \times 10^{17} \text{ g s}^{-1}$ for the two
37 wind components observed in the stable state, and about 2.5
38 and $3.9 \times 10^{16} \text{ g s}^{-1}$ in the heartbeat state.

39 From modeling the unstable accretion disk, which is par-
40 tially stabilized by the wind ejected on the cost of the locally
41 dissipated energy flux, we obtained mass outflow rates consis-
42 tent with those observed in the data. The wind outflowing as a
43 first, faster component, is probably launched from the accretion
44 disk between ~ 380 and $4700\text{--}5900 R_g$, while the second, slower
45 component probably originates between ~ 950 and $4200\text{--}4900$
46 Schwarzschild radii. The mass-loss rate \dot{M}_{wind} in these two com-
47 ponents probably is on the order of $\lesssim 4 \times 10^{16} \text{ g s}^{-1}$ if the wind
48 is weak (undetectable in our *Chandra* data) and heartbeat oscil-
49 lations are still present and on the order of $\lesssim 4 \times 10^{17} \text{ g s}^{-1}$ if the
50 wind is strong while the oscillations cease.

51 Now, the question remains whether the bound wind, ejected
52 with the velocities that are below the escape velocity for a black
53 hole with $6 M_\odot$ from the inner parts of the accretion disk, is ob-
54 servationally detectable. We suggest that this part of the flow,
55 also crucial for stabilizing the accretion disk oscillations, might
56 contribute to the hard X-ray corona. Indeed, as Fig. 2 shows,
57 there is no strong correlation between the flaring/nonflaring ac-
58 tivity and the 15–50 keV hard X-ray emission. The hard X-ray
59 emission slightly decreases during the flaring states. In the cur-
60 rent analysis, however, we cannot quantitatively describe this
61 part of the flow. The assumed time lags between the soft and
62 hard X-rays in the case when a moderate coronal flow develops




Fig. 8. Schematic geometry of the accretion flow in IGR J17091. The accretion disk dominated by radiation pressure (thin solid line) is stabilized by a quasi-static corona in its inner parts and by the unbound wind ejected at its outer radii (its velocity field is marked by arrows). The wind (we show only the first component, for clarity) is partially collimated, and its radial extension in the observed wind is somewhat narrower than at its base.

in the flaring state might be on the order of a second (Janiuk & Czerny 2005), which means that they are also beyond the scope of our current work, and it will be interesting to investigate this in the future.

In Fig. 8, we show an overall schematic geometry of the accretion flow in IGR J17091. The thin accretion disk first develops a quasi-static corona, up to a radius $r_1 \sim 380r_g$, where the outflow velocity exceeds the escape velocity from the BH gravitational potential. Above this radius, the outflow converts into the unbound wind and is ejected from the disk up to a radius $R_{1,max} \sim 4900r_g$, with a mass-loss rate that is a function of the locally dissipated energy flux in the accretion disk. Moderate mass-loss regulates the amplitude of the disk flares that are caused by the thermal-viscous instability. Strong mass-loss rate suppresses these flares completely while giving observable features in the spectra from the wind. The filling factor of this wind is consistent with the extension of the wind being similar to its launching radius, that is, $\Delta r \approx r_1$. This implies that the outflow is somewhat collimated already at the height where the spectral features are produced. In Fig. 8 only one component of the wind is shown, the other component, which has a different velocity and is also detected in the observations, has a similar geometry.

4.2. Mechanisms of launching winds from accretion disks

Models of accretion disks postulate several different ways of accelerating particles from the disk surface and of producing the unbound outflows. Winds powered by magnetic fields, radiation, or thermal driving probably operate in active galaxies (Proga 2007). Wind-launching mechanisms as a result of the radiation pressure force in their emission lines require an effective strong flux in the UV band, which means that the distance from the black hole is determined by the temperature of the flow. The temperature profile is sensitive to the black hole mass and accretion rate, but this mechanism seems implausible for the very hot accretion disks in Galactic binaries, who predominantly radiate in the soft X-ray band.

The open magnetic field lines may be responsible for the centrifugally driven outflows (Blandford & Payne 1982), and the joint action of MHD and radiation pressure forces probably

1 plays a role in luminous AGN (Proga & Kallman 2004; Everett
2 2005). For the Galactic X-ray binaries, similar mechanisms of
3 wind and jet production are commonly suggested, with the radio
4 luminosity being an observational indicator of the collimated jet
5 (Merloni et al. 2003).

6 In the microquasar GRO J1655-40, Miller et al. (2006) de-
7 tected a wind component whose absorption line properties indi-
8 cate that the magnetic field is the main driving mechanism re-
9 sponsible for the transfer of energy to the upper disk atmosphere
10 layers as well as for the outward transfer of angular momentum
11 in the accretion disk. The other two other wind-driving mecha-
12 nisms, the thermally driven wind and radiation-pressure-driven
13 wind, are excluded. The first, based on the inverse Compton
14 temperature, can only occur at very large distances of $R >$
15 $3 \times 10^6 R_{S\text{chw}}$, while the observations indicated a mean radius
16 of $200 R_{S\text{chw}}$. The second mechanism is excluded because the
17 ionization parameter for the momentum force estimated for this
18 wind is too large. Eventually, therefore, we conclude that 25%
19 of the viscously dissipated flux, if used to drive the wind against
20 the gravity force, would be enough to account for the observed
21 geometry, where the height of the wind is about $0.15 R$.

22 In our case, the thermally driven wind is also rather excluded.
23 If it was to be launched at the very large distance from the black
24 hole, then the regions of adequate temperature range could be
25 outside the disk outer edge. We estimated the outer edge location
26 of the accretion disk, based on the Roche lobe size estimate given
27 by Paczyński (1971) assuming the Main Sequence donor star
28 of $1 M_{\odot}$, and the orbital period of 4 days (Wijnands et al. 2012),
29 to be about $R_{\text{out}} \sim 4 \times 10^4$ Schwarzschild radii.

30 4.3. Viability of the radiation pressure instability model

31 The best-studied example of the radiation pressure instability in
32 action is the microquasar GRS 1915+105, which in some spec-
33 tral states exhibits cyclic X-ray outbursts that fit limited cycle os-
34 cillations of an accretion disk well. This source has been known
35 for 20 years, and only recently another microquasar of that type
36 was discovered. This is IGR J17091, the second excellent can-
37 didate source that shows radiation-pressure-driven variability on
38 observable timescales.

39 In addition, other selected Galactic X-ray binaries that were
40 discussed in more detail in Janiuk & Czerny (2011) support that
41 there are P_{rad} instabilities in their accretion disks. Nevertheless,
42 many sources accrete at a high accretion rate to even Eddington
43 rate ratios that show no limited cycle oscillations on timescales
44 adequate for the radiation pressure instability. Some stabilizing
45 mechanisms need to be considered to explain the apparent sta-
46 bility of the high accretion rate sources. One plausible mecha-
47 nisms discussed in the literature that may affect the stability of
48 accretion disks is the jet or wind outflow. Another effect that has
49 been discussed recently (Kunert-Bajraszewska et al. 2010) is the
50 potentially stabilizing effect of a companion star in a binary sys-
51 tem, or a companion galaxy, in case of quasars. Finally, a phys-
52 ically plausible possibility is viscosity propagation throughout
53 the accretion disk, as found by Janiuk & Misra (2012), which
54 suppresses the oscillations in the viscous timescale, leaving only
55 those on the thermal timescale.

56 To test the latter possibility, we used our recently developed
57 scheme of propagating viscous fluctuations in the unstable disk
58 (King et al. 2004; Lyubarskii 1997; Janiuk & Misra 2012). The
59 fluctuations, which propagate as in the Markov chain model, sta-
60 bilize the disk and lead to purely stochastic variability when only
61 the fluctuating part in the viscosity, $\alpha(r, t)$, is large. The larger the

62 fluctuating part, the more stochastic flickering is produced by the
63 disk, and finally, no heartbeat oscillations remain. Nevertheless,
64 neither of these possibilities considers mass outflow from the
65 disk and its correlation with heartbeat/no-heartbeat states, which
66 is clearly observed.

67 The luminosity outbursts for the pure radiation pressure in-
68 stability are too large with respect to the amplitudes observed in
69 IGR J17091. The most plausible scenario is that the energetic
70 wind is launched from the accretion disk and partially stabilizes
71 it, depending on its strength and mass-loss rate. We determined
72 the range of plausible wind strengths based on the timescales
73 and amplitudes of the disk flares they allow. The wind may ei-
74 ther partially stabilize the disk oscillations to produce regular
75 outbursts, but of moderate amplitudes, or even completely stabi-
76 lize the disk.

5. Conclusions

78 We studied the microquasar IGR J17091 with respect to
79 its unique properties. It is the second known source (after
80 GRS 1915+105) that shows the limited-cycle oscillations char-
81 acteristic for the thermal-viscous instability of an accretion disk.
82 For the physical parameters of the source that are known from
83 observations, albeit with some uncertainties (its black hole mass
84 and Eddington ratio), and assuming the still-used phenomeno-
85 logical α -viscosity prescription of the Shakura-Sunyaev model,
86 the hydrodynamical evolution of the disk shows regular lumi-
87 nosity oscillations. Nevertheless, for the required timescales and
88 amplitudes of these observed oscillations, as well as to suppress
89 them further whenever the source enters its quasi-static mode,
90 we need to account for an additional physical mechanism to
91 remove some fraction of the viscously dissipated flux and par-
92 tially (or completely) stabilize the disk. The best process in this
93 context is launching of the outflow from the accretion disk, as
94 shown not only theoretically, but also motivated by the indepen-
95 dent spectroscopic observations of this source.

96 We demonstrated that the wind strength extracted from the
97 data agrees well with our theoretical scenario. Even assum-
98 ing that part of the wind is not observable because it is to-
99 tally ionized, we verified that in any case the mass outflow rate
100 ($3\text{--}4.7 \times 10^{17} \text{ g s}^{-1}$) of the observed wind in the no-heartbeat
101 state is higher than the rate needed in our scenario to totally sta-
102 bilize the heartbeat. Moreover, the lack of any lines detected in
103 the *Chandra* spectra suggests for the heartbeat state an upper
104 limit for the mass outflow rate that is compatible with the rate
105 obtained by our model to obtain a flare amplitude that agrees
106 with the observed amplitudes ($2.5\text{--}4 \times 10^{16} \text{ g s}^{-1}$).

Our final conclusions are as follows:

- The heartbeat oscillations of the X-ray luminosity of IGR J17091 detected during its outburst in 2011 are attributed to the radiation pressure instability of the accretion disk.
- The outflow launched from the disk at the cost of part of the locally dissipated energy flux is a plausible mechanism to regulate the amplitude of these oscillations.
- The strong outflow may stabilize the disk and completely suppress the heartbeat.
- The observed properties of the wind detected from IGR J17091 in the state without the heartbeats allowed us to constrain the the mass-loss rate at large distances in the disk.
- The MHD-driven wind from the accretion disk, used in our model to stabilize the disk, is strong in the innermost parts.

1 In the spectroscopic observations, it may not always be de-
 2 tectable because of its high ionization state or because the
 3 velocities required very close to the black hole are below the
 4 escape velocity; in that case, the MHD mechanism instead
 5 drives the formation of a quasi-static bound corona above
 6 the disk. However, we demonstrated that the observed winds
 7 agree well with the scenario described by our model.

8 *Acknowledgements.* We thank Bozena Czerny, Daniel Proga, and Agata
 9 Rozanska for helpful discussions. This work was supported in part by the grant
 10 DEC-2012/05/E/ST9/03914 from the Polish National Science Center. We also
 11 acknowledge support from the COST Action MP0905, “Black Holes in a Violent
 12 Universe”.

13 References

14 Abramowicz, M. A., Czerny, B., Lasota, J. P., & Szuszkiewicz, E. 1988, *ApJ*,
 15 332, 646
 16 Altamirano, D., & Belloni, T. 2012, *ApJ*, 747, L4
 17 Altamirano, D., Belloni, T., Krimm, H., et al. 2011a, *ATel*, 3230, 1
 18 Altamirano, D., Belloni, T., Linares, M., et al. 2011b, *ApJ*, 742, L17
 19 Altamirano, D., Linares, M., van der Klis, M., et al. 2011c, *ATel*, 3225, 1
 20 Belloni, T., Klein-Wolt, M., Méndez, M., van der Klis, M., & van Paradijs, J.
 21 2000, *A&A*, 355, 271
 22 Blandford, R. D., & Payne, D. G. 1982, *MNRAS*, 199, 883
 23 Canizares, C. R., Davis, J. E., Dewey, D., et al. 2005, *PASP*, 117, 1144
 24 Capitanio, F., Ubertini, P., Bazzano, A., et al. 2005, *ApJ*, 622, 503
 25 Capitanio, F., Giroletti, M., Molina, M., et al. 2009, *ApJ*, 690, 1621
 26 Capitanio, F., Del Santo, M., Bozzo, E., et al. 2012, *MNRAS*, 422, 3130
 27 Cash, W. 1976, *A&A*, 52, 307
 28 Corbel, S., & Tzioumis, T. 2011, *ATel*, 3246, 1
 29 Corbel, S., Rodriguez, J., Tzioumis, T., & Tomsick, J. 2011, *ATel*, 3167, 1
 30 Czerny, B., Siemiginowska, A., Janiuk, A., Nikiel-Wroczyński, B., & Stawarz,
 31 Ł. 2009, *ApJ*, 698, 840
 32 Done, C., Wardziński, G., & Gierliński, M. 2004, *MNRAS*, 349, 393
 33 Everett, J. E. 2005, *ApJ*, 631, 689
 34 Ferland, G. J., Porter, R. L., van Hoof, P. A. M., et al. 2013, *Rev. Mex. Astron.*
 35 *Astrofis.*, 49, 137
 36 Fruscione, A., McDowell, J. C., Allen, G. E., et al. 2006, in *SPIE Conf. Ser.*,
 37 6270, 62701V
 38 Garmire, G. P., Bautz, M. W., Ford, P. G., Nousek, J. A., & Ricker, Jr., G. R.
 39 2003, in *X-Ray and Gamma-Ray Telescopes and Instruments for Astronomy.*,
 40 eds. J. E. Truemper, & H. D. Tananbaum, *SPIE Conf. Ser.*, 4851, 28
 41 in ’t Zand, J. J. M., Heise, J., Lowes, P., & Ubertini, P. 2003, *ATel*, 160, 1

Janiuk, A., & Czerny, B. 2005, *MNRAS*, 356, 205 42
 Janiuk, A., & Czerny, B. 2011, *MNRAS*, 414, 2186 43
 Janiuk, A., & Misra, R. 2012, *A&A*, 540, A114 44
 Janiuk, A., Czerny, B., & Siemiginowska, A. 2000, *ApJ*, 542, L33 45
 Janiuk, A., Czerny, B., & Siemiginowska, A. 2002, *ApJ*, 576, 908 46
 Jourdain, E., Götz, D., Westergaard, N. J., Natalucci, L., & Roques, J. P. 2008,
 in *Proc. of the 7th INTEGRAL Workshop*, Copenhagen, Denmark 47
 Kennea, J. A., & Capitanio, F. 2007, *ATel*, 1140, 1 48
 King, A. R., Pringle, J. E., West, R. G., & Livio, M. 2004, *MNRAS*, 348, 111 49
 King, A. L., Miller, J. M., Raymond, J., et al. 2012, *ApJ*, 746, L20 50
 Kunert-Bajraszewska, M., Janiuk, A., Gawroński, M. P., & Siemiginowska, A.
 2010, *ApJ*, 718, 1345 51
 Kuulkers, E., Lutovinov, A., Parmar, A., et al. 2003, *ATel*, 149, 1 52
 Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141 53
 Lutovinov, A. A., & Revnivtsev, M. G. 2003, *Astron. Lett.*, 29, 719 54
 Lutovinov, A., Revnivtsev, M., Molokov, S., & Sunyaev, R. 2005, *A&A*, 430, 997 55
 Lyubarskii, Y. E. 1997, *MNRAS*, 292, 679 56
 Merloni, A., Heinz, S., & di Matteo, T. 2003, *MNRAS*, 345, 1057 57
 Miller, J. M., Raymond, J., Fabian, A., et al. 2006, *Nature*, 441, 953 58
 Muchotrzeb, B., & Paczynski, B. 1982, *Acta Astron.*, 32, 1 59
 Nayakshin, S., Rappaport, S., & Melia, F. 2000, *ApJ*, 535, 798 60
 Neilsen, J., Petschek, A. J., & Lee, J. C. 2012, *MNRAS*, 421, 502 61
 Paczyński, B. 1971, *ARA&A*, 9, 183 62
 Pahari, M., Yadav, J. S., & Bhattacharyya, S. 2014, *ApJ*, 783, 141 63
 Pandey, M., Manchanda, R. K., Rao, A. P., Durouchoux, P., & Ishwara-Chandra.
 2006, *A&A*, 446, 471 64
 Piconcelli, E., Jimenez-Bailón, E., Guainazzi, M., et al. 2004, *MNRAS*, 351, 161 65
 Ponti, G., Fender, R. P., Begelman, M. C., et al. 2012, *MNRAS*, 422, L11 66
 Proga, D. 2007, in *The Central Engine of Active Galactic Nuclei*, eds. L. C. Ho,
 & J.-W. Wang, *ASP Conf. Ser.*, 373, 267 67
 Proga, D., & Kallman, T. R. 2004, *ApJ*, 616, 688 68
 Rao, A., & Vadawale, S. V. 2012, *ApJ*, 757, L12 69
 Rebusco, P., Moskalik, P., Kluźniak, W., & Abramowicz, M. A. 2012, *A&A*,
 540, L4 70
 Reis, R. C., Miller, J. M., King, A. L., & Reynolds, M. T. 2012, *ATel*, 4382, 1 71
 Revnivtsev, M., Gilfanov, M., Churazov, E., & Sunyaev, R. 2003, *ATel*, 150, 1 72
 Rodriguez, J., Corbel, S., Caballero, I., et al. 2011a, *A&A*, 533, L4 73
 Rodriguez, J., Corbel, S., Tomsick, J. A., Paizis, A., & Kuulkers, E. 2011b, *ATel*,
 3168, 1 74
 Rupen, M. P., Mioduszewski, A. J., & Dhawan, V. 2003, *ATel*, 152, 1 75
 Shakura, N. I., & Sunyaev, R. A. 1973, *A&A*, 24, 337 76
 Shaposhnikov, N. 2011, *ATel*, 3179, 1 77
 Strüder, L., Briel, U., Dennerl, K., et al. 2001, *A&A*, 365, L18 78
 Torres, M. A. P., Jonker, P. G., Steeghs, D., & Mulchaey, J. S. 2011, *ATel*, 3150,
 1 79
 Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131 80
 Wijnands, R., Yang, Y. J., & Altamirano, D. 2012, *MNRAS*, 422, L91 81
 Yu, W., & Yan, Z. 2009, *ApJ*, 701, 1940 82
 83
 84
 85
 86
 87
 88
 89